



Idaho

Live births, abortions, and pregnancies


Live birth rates, 1982–1992



Live birth rates,* 1991



Pregnancy rates, 1990



National live birth rate ranking, 1990


Age group		
15-17	18-19	15-19
38th highest	30th highest	34th highest

*Missing bar indicates fewer than 10,000 females or fewer than 20 births. This graph refers to females aged 15–19 years.




Sexually transmitted diseases Idaho

Gonorrhea rates,* 1982–1992



Gonorrhea rates, 1992



National ranking,[†] 1992

Males – 47th highest Females – 46th highest


*Data were unavailable before 1986.

†National ranking refers to males and females aged 10–24 years.




Chlamydia cases and rates,* 1988–1992


Cases




Rates




Cases



Rates



Acquired immunodeficiency syndrome (AIDS) cumulative cases,† 1981–1992



*Chlamydia cases and rates refer to males and females of all ages.


†Missing bar indicates fewer than six cumulative cases.



Related risk behaviors,* 1993


Idaho

Had sexual intercourse (lifetime)




Question not included in survey.

Had sexual intercourse with four or more people (lifetime)




Question not included in survey.

Had sexual intercourse during the past three months (currently sexually active)




Question not included in survey.

Used a condom during last sexual intercourse (among currently sexually active)




Question not included in survey.

Used birth control pills during last sexual intercourse (among currently sexually active)



Question not included in survey.

Injected illegal drugs (lifetime)



*Related risk behaviors refer to high school students.
Sources: CDC—NCHS and NCCDPHP (birth, abortion, and pregnancy data), NCPS (gonorrhea and chlamydia data), NCID (AIDS data); 1993 Idaho Youth Risk Behavior Survey (risk behavior data).

